

Kerala Agricultural University, Vellanikkara, Thrissur Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in & kauagmet@yahoo.co.in







ERNAKULAM DISTRICT

AAS 41/2018(82), Friday

12.10.2018

The Agricultural advisory of this bulletin is not applicable in flood-affected areas.

CURRENT SYNOPTIC SITUATION: A Deep Depression lies over Odisha centred near Lat 20.6° N/Long. 84.9° E. A severe cyclonic storm (LUBAN) lies over west Central Arabian sea near Lat 14.7° N/Long. 57.4° E. A trough runs from the deep depression to Kerala across Coastal Andhra Pradesh & Tamil Nadu and extends upto 3.1km Above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	13.10.2018	14.10.2018	15.10.2018	16.10.2018	17.10.2018
Rainfall (mm)	2	8	4	0	12
Maximum temperature (°C)	32.0	31.8	31.9	31.6	31.8
Minimum temperature (°C)	24.6	24.4	24.2	24.2	24.2
Total cloud cover (octa)	6	6	6	6	6
Maximum Relative Humidity (%)	90	90	90	90	90
Minimum Relative Humidity (%)	65	65	65	65	65
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	320	320	320	320	320

As per IMD forecast, light rainfall in isolated places is expected for coming days in Ernakulam district.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Crop stage	Pest/ Disease	Advisories		
Paddy	Harvesting		First paddy crop is in Harvesting stage. Land preparation for		
	stage	-	second paddy crop can be started.		
Coconut	_	Rhinoceros	After through cleaning of the crown region, apply cartap hydrochloride 4G @ 20 g mixed with fine sand 200 g to the inner most leaf axils surrounding the spindle leaf or a mixture of fipronil 0.3		
		beetle	G @ 20 g and fine sand @ 200 g to the inner most leaf axils. Tthis application has to be repeated thrice a year. (January, May and September/ October) for better results.		
Banana	-	Spodoptera	Destroy the affected leafs along with the spodoptra. For severe attack spray quinalphos 2ml per one litre of water or chlorantraniliprole 3 ml/10 litre of water.		
Animal	• Animal should not be left to graze under hot sun from morning 10.00 AM to evening 3.00 PM.				
Husbandry	Concentrate feed should be sun dried regularly to avoid mould growth				
	Deworm the animals and birds and vaccinate them against infectious diseases.				
	(CAADECCS, KVASU, Mannuthy)				