


Agromet Advisory Bulletin for the District, Kasaragod

(Valid from 17.09.2025 to 21.09.2025)





Bulletin Number: Pilicode/ Ksd – 74/2025	Date: 16/09/2025
--	------------------

A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
5.4	30.0 - 30.2	23.0 - 24.6	79 – 96	00 - 01

B. Weather forecast for next five days

Parameters	17-09-2025	18-09-2025	19-09-2025	20-09-2025	21-09-2025
Average Rainfall, mm	2	10	4	0.1	0.1
Max. Temp, °C	31	31	31	31	31
Min. Temp,°C	24	24	24	24	24
Max. Relative Humidity, %	86	86	86	86	86
Min. Relative Humidity, %	61	61	61	61	61
Wind speed,km/h	3	3	3	4	3
Wind direction, degrees	320	340	320	290	320
Total cloud cover, octa	8	8	8	7	8

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories	
General	Light to Moderate Rainfall** The sky will be partially or fully cloudy. High relative humidity will be experienced. The atmospheric temperature will be high.			
Condition				
Weather	There will be heavy ra	There will be heavy rainfalls on September 17 in isolated places.		
warning	There will be light to moderate rainfalls from September 18 to 20.			
Impacts	Damaging and losses or deteriorating quality of the open drying agricultural produces like copra, rubber, cashew nuts, seeds, food materials for processing etc			
	Damages and losses may occur for crops having slender stems like banana and vegetables. Breaking branches and or trees may occur.			
	High temperature during daytime may cause high rate of evaporation of water from the soil.			
General Recommendati	Do not attempt to dry the products like rice, rubber, copra, pulses, vegetable seeds, etc directly under sun.			
ons	Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc. Clean the drainage channels in crop lowlands to enable proper drainage of excess water in case of exigencies if any.			
	Do not shelter the cattle in houses with weak structures.			
	Keep animal feeds, seeds, fertilizers etc. in termite free and moist free rooms on elevated platforms coated with wooden panels.			
	Provide mulching to crops to reduce waterloss from soil.			

Coconut	All stages	Bud rot	Detection of disease at its early stage will help to adopt efficient corrective measures. Cut and remove the affected tissues from the crown and apply Bordeaux paste. After that cover the cut surface with polythene sheets to protect it from rain falls until new leaf emerges. Burn the removed tissues immediately. As a prophylactic measure spray 1% Bordeaux mixture into the axils of top leaves of the surrounding palms
Coconut	All stages	Yellowing of coconut	Incorporate lime or dolomite @ 2 kg/palm to the basins of the palms. After two weeks, apply borax @ 200g/palm and zinc sulphate @ 100g/palm and magnesium sulphate @500g/palm and potash @ 2kg/palm
Areca nut	All stages	Mahali disease	Apply Bordeaux mixture (1%) to the bunch and leaves.
Black pepper	Bearing stage	Pollu beetle/disease	As a pyrolytic measure spray spray 1% Bordeaux mixture mixed with quinalphos (2ml/litre of Bordeaux mixture). If attack is seen spray solution containing hexaconazole (5ml) + cyhalothrin (3ml) per 5 litres of water.
Cashew	Pre-bearing stages	Tea mosquito bug	Prophylactic measures: 1) The fungal biological control agent, Beavaria bassiana can be sprayed @ 20g/L Or 2) Spray 1% Bordeaux mixture mixed with quinalphos (2ml/litre of Bordeaux mixture)

1	
Banana	Various stages of growth



There should not be any water stagnation in the basins of the crop.

As a prophylactic measure spray Pseudomonas (@ 20g/litre) at fortnight intervals.

If disease appears:-Cut and burn all affected leaves. Spray Carbendazim and Mancozeb (@ 2ml per litre) alternatively at fortnight intervals

** Warning colour codes of rainfall (for disaster management)

Warning (Take actions) Alert (Be prepared) Watch (Be u	
	e updated) No warning (No actions)

Sd/-Nodal Officer, GKMS Project, RARS Pilicode