





## Agromet Advisory Bulletin for PALAKKAD District (Valid from 09.07.2025 to 13.07.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

## AAS Bulletin No: 27/2025, (54) Tuesday

Dated 08.07.2025

## A. Medium range weather forecast of IMD for next five days

Weather parameters	09.07.2025	10.07.2025	11.07.2025	12.07.2025	13.07.2025
Rainfall (mm)	10	6	8	10	5
Maximum temperature (°C)	29	29	29	29	29
Minimum temperature (°C)	23	23	23	23	23
Maximum Relative Humidity (%)	95	95	95	95	95
Minimum Relative Humidity (%)	85	85	85	85	85
Wind speed (kmph)	5	5	5	5	5
Wind direction (deg)	250	270	250	250	250
Total cloud cover (octa)	7	6	6	6	6

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of light rainfall in the district and moderate rainfall in isolated places of the district. Strong winds speed reaching likely to prevail occasionally over Kerala from 08th to 10th July 2025. Above normal rainfall is expected in Kerala.

## **B.** Agrometeorological Advisories for crop protection & management

General and Agrometeorological Advisories for Palakkad District

General Advisory: -Make proper drainage facilities in the crop fields. Give propping to the plants and make it stronger. Harvest the matured crop fields as early as possible. Make the land free from weeds, dried leaves and diseased plants to avoid the attack of pests and disease.

Сгор	Pest/ Disease, Crop Management	Agromet Advisory		
Rice (virippu)	Thrips	To control thrips in rice, spray 2 gram of Thiomethoxam in 10 litre of water. Pesticides should not be applied during rainy period.		
Coconut	Stem bleeding	As it is raining, there is a chance for the development of stem bleeding disease in coconut palms. As a precaution, apply 5 kg neem cake in the basins of the palms. Also, apply 50g each trichoderma mixed along with organic manures in the basins.		
Pepper	Quick wilt	Drench the soil with 0.2% copper oxy chloride and spray the aerial part with 1% Bordeaux mixture as a prophylactic measure		
Animal Husbandry: Sundry the cattle feed frequently to avoid fungal growth. Give foot and mouth disease (FMD) vaccination to the animal. Don't allow the animals go to water logged areas in order to avoid anthrax and foot lesions. (CAADECCS, KVASU, Mannuthy)				