



## Agromet Advisory Bulletin for Thrissur District

(Valid from 15.06.2024 to 19.06.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara  
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

**AAS Bulletin No: 24/2024, (48) Friday**

**Dated 14.06.2024**

Observed weather during last week (Dated 08 <sup>th</sup> June to 14 <sup>th</sup> June 2024)							Weather Parameters		Weather Forecast Valid for 15 <sup>th</sup> June to 19 <sup>th</sup> June 2024				
08/06	09/06	10/06	11/06	12/06	13/06	14/06	Date		15/06	16/06	17/06	18/06	19/06
6.7	106.1	11.0	12.6	1.0	1.1	0.0	Rainfall (mm)		2	4	7	18	10
31.8	31.0	31.7	31.5	31.0	29.4	32.2	T <sub>max</sub> ( <sup>o</sup> C)		32	32	32	32	32
24.9	23.9	24.5	24.5	24.1	24.6	24.6	T <sub>min</sub> ( <sup>o</sup> C)		25	25	25	25	25
Cloudy	Cloudy	Overcast	Cloudy	Overcast	Cloudy	Partially cloudy	AM	Cloud cover	Overcast	Overcast	Overcast	Overcast	Overcast
Overcast	Cloudy	Cloudy	Cloudy	Overcast	Partially cloudy	PM							
95	97	97	97	93	97	95	RH-I (%)		90	90	90	90	90
81	74	80	91	83	71		RH-II (%)		77	77	77	77	77
1.9	1.4	2.5	1.6	1.2	0.3	1.8	Wind Speed (km/hr.)		3	4	2	2	2
C	C	C	C	C	C	C	AM	Wind direction	SW	WSW	WSW	SW	WSW
SSW	SSW	WSW	C	C	WSW	PM							
Rainfall (mm) in last week				Rainfall (mm) from 01/01/2024 to till dated			Rainfall (mm) from 01/06/2024 to today						
138.5				973.7			342.9						

**Weather summary / Alert:** As per the forecast of IMD, there is a chance of Moderate rainfall in isolated places of the district. Below normal rainfall is expected in Kerala.

### General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops before monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports for the bananas. Make the vegetable pandals stronger. Avoid standing in open areas to protect farmers and livestock from lightning and thunder.

Crop		Crop/ Animal Advisory
Rice (virippu)	Sowing/Transplantig	Time of land preparation in paddy fields, 140 kg of lime can be applied per acre. During rainy period, there is a chance of sheath blight, sheath rot, leaf spot disease etc. As a prophylactic measure, either soak one kg of paddy seeds or dip root of seedlings in a solution of Pseudomonas culture @10gram of Pseudomonas culture per one litre of water for 30 minutes.
Banana	Rhizome rot disease	During rainy season, there is a chance of Rhizome rot in banana. Proper drainage facilities have to be ensured. As a precaution, drench the basin with three litre of pseudomonas solution @ 20g per one liter of water.
Nutmeg	Organic fertilization	One year old plants can be given 10 kg of manure or compost per plant.
Pepper	Organic fertilization	At the beginning of the rainy season the first dose of fertilizer can be applied to pepper plants. A basin of 10 to 15 cm depth and 30 to 40 cm. diameter should be taken. 10 kg of manure or compost can be applied per plant and add some soil.

**Animal Husbandry:** - Deworm the animals after dung examination. Take precautions to avoid ecto parasitic infestation. Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis..